

## Comparative Analysis of Pulmonary Function Tests (PFTs) In Athletes: Diagnostic and Clinical Relevance

Janu S. Chandak and Peryben Y. Patel\*

*Department of Physiotherapy, Subhadraben Sureshchandra Institute of Physiotherapy,  
Faculty of Health and Allied Sciences, Ganpat University, Ganpat Vidyanagar, Dist. Mehsana, Gujarat, India.*

### Abstract

Pulmonary Function Tests (PFTs) play a critical role in assessing lung health and performance, particularly in athletes exposed to unique respiratory challenges due to intense training and environmental stressors. This review synthesizes recent findings on the application of PFTs in athletic populations. It highlights variations in test outcomes based on sport type, gender, training intensity, and environmental exposure. Furthermore, it explores how PFTs aid in diagnosing exercise-induced bronchoconstriction (EIB), asthma, and other respiratory conditions, while discussing emerging diagnostic tools such as specific airway conductance (SGaw), eucapnic voluntary hyperpnoea (EVH), and oscillometry. The paper also outlines clinical implications including treatment monitoring, athlete screening, and anti-doping compliance. Collectively, the findings reinforce the value of PFTs in enhancing respiratory care and optimizing athletic performance.

**Keywords:-** Pulmonary Function Tests, lung health, Athletic populations, exercise-induced bronchoconstriction

### 1. Introduction:

Athletes frequently exhibit lung function profiles that deviate from general population norms due to chronic high-intensity training and specific environmental exposures.(Crapo et al.,2025) These differences can signify either adaptive physiological changes or underlying respiratory disorders.(Stensrud et al., 2010) For instance, elite athletes may experience symptoms such as dyspnea or wheezing, which may be attributed to conditions like asthma or exercise-induced bronchoconstriction (EIB), or merely reflect normal adaptation.(Smith et al., 2002) As subjective symptoms are often unreliable, PFTs provide objective and measurable data critical to evaluating respiratory health in athletes.(Parsons et al., 2005) This review examines the diagnostic utility of Pulmonary Function Tests (PFTs) across diverse athlete demographics and sports disciplines.(Hiemstra et al., 2018)

### 2. Overview of Pulmonary Function Tests

PFTs assess various aspects of lung function, including airflow, volume, and gas exchange. Spirometry, the most widely used test, evaluates key parameters such as forced expiratory volume in one second (FEV1), forced vital capacity (FVC), and the FEV1/FVC ratio. Additional

techniques include:

- **Plethysmography:** Measures static lung volumes and airway resistance.(Stensrud et al., 2010, Smith et a., 2002)
- **DLCO (Diffusing Capacity of Lung for Carbon Monoxide) :** Assesses the lungs' ability to transfer gases into the bloodstream.(Smith et a., 2002, Parsons et al., 2005)
- **SGaw (specific airway conductance):** Indicates airway openness and is sensitive to early obstruction.(Parsons et al., 2005)
- **EVH (Eucapnic Voluntary Hyperpnoea) and Broncho provocation tests:** Particularly valuable in diagnosing EIB when spirometry is inconclusive.(Hiemstra et al., 2018)

Environmental conditions, recent physical exertion, time of day, and baseline lung capacity all influence results and should be carefully controlled or accounted for in testing protocols.

\*Corresponding Author  
E-mail address: perry Patel101@gmail.com




Figure – 1 Pulmonary Function Tests (PFT) – Techniques and influencing factors

### 3. How PFTs Help Identify Respiratory Conditions

#### 3.1 Exercise-Induced Bronchoconstriction (EIB)

EIB is characterized by a temporary narrowing of the airways following exercise and is especially prevalent among endurance athletes. Diagnosis typically involves identifying a 10–15% decrease in FEV1 after exercise. Since symptoms may not correlate with actual bronchoconstriction, post-exercise spirometry and EVH are more reliable diagnostic tools. (Dempsey et al., 1999, Madigan et al., 2002)

#### 3.2 Distinguishing Asthma from EIB

Asthma is a chronic inflammatory condition, whereas EIB is a transient phenomenon. Baseline spirometry in asthma often shows obstruction reversible with bronchodilators. (Andreotta et al., 2018) In contrast, athletes with isolated EIB may have normal baseline values but exhibit airway narrowing only after exercise or during challenge testing. Differentiating the two is essential for effective management and treatment planning. (Boulet, 2012)

#### 3.3 Obstructive vs. Restrictive Patterns

Endurance athletes, such as long-distance runners, may exhibit obstructive patterns due to airway remodelling. Strength athletes may show restrictive patterns due to increased thoracic musculature. Comparative studies indicate endurance athletes typically have higher ventilatory capacity and gas diffusion efficiency, while strength athletes demonstrate greater inspiratory pressure but reduced airflow rates. (Mahler et al., 1993, Randolph et al., 2013)

#### 3.4 Environmental Factors

Environmental exposures significantly impact respiratory health. Cold air, airborne pollutants, allergens, and chlorinated pool environments can provoke airway irritation. Skiers and hockey players often experience EIB due to cold air inhalation, while swimmers may develop chlorine-induced airway inflammation. Testing under

sport-specific environmental conditions improves diagnostic accuracy. (Kitzmüller et al., 2011, Tiller et al., 2019)

#### 3.5 Use of Advanced Testing

Advanced diagnostic modalities are becoming more prevalent in sports medicine:

- SGaw (specific airway conductance): More sensitive than spirometry in detecting airway obstruction.
- Oscillometry: Provides information on small airway function with minimal effort.
- Flow-volume loops: Useful for identifying inspiratory limitations and extra thoracic obstruction. These tools allow for more nuanced and comprehensive assessment, especially in elite and symptomatic athletes. (Randolph, 2013)




Figure 2 – Respiratory Conditions In athletes

### 4. Comparison by Sport, Gender, Age, and Intensity

Athletic discipline significantly influences lung function profiles. Endurance sports like swimming and cycling typically enhance lung volume and efficiency, while team sports involving intermittent effort may result in more variable patterns. (Krafczyk et al., 2011)

Gender Differences: Female athletes generally have smaller airway diameters and lung volumes, increasing

their susceptibility to respiratory symptoms. Hormonal fluctuations may further influence airway reactivity.(Mckenzie et al.,2011)

**Age-Related Variability:** Adolescent athletes may be underdiagnosed due to incomplete lung development. Older athletes often experience natural declines in lung function, which must be considered in test interpretation.(Constantinou et al., 2010)

**Training Altitude:** High-altitude training temporarily enhances oxygen diffusion but may reduce lung volumes over time. These variables must be contextualized in performance and diagnostic evaluations.(Wilcox et al., 200




Figure 3 – Comparison by Various outcomes




Figure 4 – Factors Influencing Lung Function In athletes

## 5. Challenges and Limitations

While PFTs are indispensable tools, they come with inherent limitations:

- **Environmental Simulation:** Laboratory conditions may not mimic real sport scenarios, resulting in false negatives.(Storms et al., 1997)
- **Resource Limitations:** Broncho provocation tests require specialized settings and personnel.(Ludyga et al., 2022)
- **Reference Standards:** There is a lack of sport-specific normative values for interpreting results.

Suggested strategies include field-based portable spirometry, athlete-specific baseline comparisons, and the use of complementary tools such as symptom checklists, training logs, and inflammation biomarkers (e.g., exhaled nitric oxide). A multidisciplinary team ensures

comprehensive assessment and appropriate intervention. (Storms et al., 1997, Ludyga et al., 2022)

## 6. Clinical Use and Athlete Care

Baseline PFTs are beneficial for early detection and tracking of respiratory issues. They serve as important references for evaluating treatment outcomes or illness recovery. Personalized management plans may include tailored medications, warm-up protocols, or environmental adjustments. Repeated testing helps monitor therapy efficacy. (Chan et al., 2020)

PFTs also play a critical role in ensuring athletes' compliance with anti-doping regulations. Medications such as inhaled corticosteroids or beta-agonists are permitted under documented medical need. Accurate diagnostic data provides essential justification and safeguards athletes' health and careers.(Cox et al., 2014)

## 7. Future Developments

Advancements in technology promise transformative improvements in respiratory monitoring:

- Wearable spirometry devices enable real-time lung function tracking during training.
- Artificial intelligence supports automated interpretation and trend analysis.

More research is needed to address underrepresented groups, including female, para-athletes, and those in extreme environments. Longitudinal studies on post-career respiratory health may inform future prevention and care models.(Ralalage et al., 2023, Needham et al., 2022, Wu et al., 2025)

## 8. Conclusion

Pulmonary Function Tests are foundational in the assessment and management of respiratory health in athletes. They enable accurate diagnosis, inform treatment strategies, and support optimal performance. As sport science evolves, integrating advanced diagnostics and individualized approaches will be key to safeguarding and enhancing athletic health.

## References

- Crapo, R. O., Casaburi, R., Coates, A. L., Enright, P. L., Hankinson, J. L., Irvin, C. G., MacIntyre, N. R., McKay, R. T., Wanger, J. S., Anderson, S. D., Cockcroft, D. W., Fish, J. E., & Sterk, P. J. (2000). Guidelines for Methacholine and Exercise Challenge Testing—1999. *American Journal of Respiratory and Critical Care Medicine*, 161(1), 309–329. <https://doi.org/10.1164/ajrccm.161.1.ats11-99>
- Stensrud, S., Myklebust, G., Kristianslund, E., Bahr, R., & Krosshaug, T. (2010). Correlation between two-dimensional video analysis and subjective assessment in evaluating knee control among elite female team handball players. *British Journal of Sports Medicine*, 45(7), 589–595. <https://doi.org/10.1136/bjism.2010.078287>
- Smith, D. J., Norris, S. R., & Hogg, J. M. (2002). Performance evaluation of swimmers. *Sports Medicine*, 32(9), 539–554. <https://doi.org/10.2165/00007256-200232090-00001>
- Parsons, J. P., Marsh, C. B., & Mastronarde, J. G. (2005). WERNICKE'S ENCEPHALOPATHY IN A PATIENT AFTER GASTRIC BYPASS SURGERY. *CHEST Journal*, 128(4), 453S. <https://doi.org/10.1378/chest.128.6.3966>
- Hiemstra, J. A., Veteto, A. B., Lambert, M. D., Olver, T. D., Ferguson, B. S., McDonald, K. S., Emter, C. A., & Domeier, T. L. (2018). Chronic low-intensity exercise attenuates cardiomyocyte contractile dysfunction and impaired adrenergic responsiveness in aortic-banded mini-swine. *Journal of Applied Physiology*, 124(4), 1034–1044. <https://doi.org/10.1152/jappphysiol.00840.2017>
- Dempsey, J. A., & Wagner, P. D. (1999). Exercise-induced arterial hypoxemia. *Journal of Applied Physiology*, 87(6), 1997–2006. <https://doi.org/10.1152/jappl.1999.87.6.1997>
- Madigan, M. L., & Pidcoe, P. E. (2002). A muscle temperature compensation technique for EMG fatigue measures. *Medicine & Science in Sports & Exercise*, 34(5), 780–784. <https://doi.org/10.1097/00005768-200205000-00009>
- Andreotta, P. W., Arold, S., Kenyon, J., Spicer, D., Woodman, P., Berry, E., Brogan, T., Kong, S., Okerholm, P., Russell, V., Clarke, R. W., & Hava, D. L. (2018). Inhaled calcium salts inhibit tobacco smoke-induced inflammation by modulating expression of chemokines and cytokines. *Pulmonary Pharmacology & Therapeutics*, 53, 86–99. <https://doi.org/10.1016/j.pupt.2018.10.004>
- Boulet, L.-P. (2012). Cough and upper airway disorders in elite athletes: a critical review. *British Journal of Sports Medicine*, 46(6), 417–421. <https://doi.org/10.1136/bjsports-2011-090812>
- Mahler, D. A. (1993). Exercise-induced asthma. *Medicine and Science in Sports and Exercise*, 25(5), 554–561.
- Randolph, C. (2013). Pediatric Exercise-Induced Bronchoconstriction: Contemporary Developments in Epidemiology, Pathogenesis, Presentation, Diagnosis, and Therapy. *Current Allergy and Asthma Reports*, 13(6), 662–671. <https://doi.org/10.1007/s11882-013-0380-x>
- Kitzmüller, C., Nagl, B., Deifl, S., Walterskirchen, C., Jahn-Schmid, B., Zlabinger, G. J., & Bohle, B. (2011). Human blood basophils do not act as antigen-presenting cells for the major birch pollen allergen Bet v 1. *Allergy*, 67(5), 593–600. <https://doi.org/10.1111/j.1398-9995.2011.02764.x>
- Tiller, N. B. (2019). Pulmonary and Respiratory Muscle Function in Response to Marathon and Ultra-Marathon Running: A Review. *Sports Medicine*, 49(7), 1031–1041. <https://doi.org/10.1007/s40279-019-01105-w>
- Eisenmann, J. C., & Malina, R. M. (2002). Growth status and estimated growth rate of young distance runners. *International Journal of Sports Medicine*, 23(3), 168–173. <https://doi.org/10.1055/s-2002-23174>
- Krafczyk, M. A., & Asplund, C. A. (2011). Exercise-induced bronchoconstriction: diagnosis and management. *American Family Physician*, 84(4), 427–434.
- Mckenzie, D. C., & Fitch, K. (2011). The Asthmatic Athlete: Inhaled Beta-2 Agonists, Sport Performance, and Doping. *Clinical Journal of Sport Medicine*, 21, 46-50. <https://doi.org/10.1097/IAE.0b013e318203c0ef>
- Constantinou, D. (2010). Exercise-induced bronchoconstriction: Current update and implications for treating athletes. *Current Allergy and Clinical Immunology*, 23, 64–70.
- Wilcox, P., Road, J., & Altalag, A. (2009). *Pulmonary Function Tests in Clinical Practice*. Springer London. <https://doi.org/10.1007/978-1-84882-231-3>
- Storms, W. W., & Joyner, D. M. (1997). Update on Exercise-Induced Asthma: A Report of the Olympic Exercise Asthma Summit Conference. *The Physician and Sportsmedicine*, 25(3), 45–55. <https://doi.org/10.1080/00913847.1997.11440200>
- Ludyga, S., Hanke, M., Leuenberger, R., Bruggisser, F., Pühse, U., Gerber, M., Lemola, S., Capone-Mori, A., Keutler, C., Brotzmann, M., & Weber, P. (2022). Martial Arts and Cognitive Control in Children with Attention-Deficit Hyperactivity Disorder and Children Born Very Preterm: A Combined Analysis of Two Randomized Controlled Trials. *Medicine & Science in Sports & Exercise*, 55(5), 777–786. <https://doi.org/10.1249/mss.0000000000003110>

- Chan, E. S., Dinakar, C., Gonzales-Reyes, E., Green, T. D., Gupta, R., Jones, D., Wang, J., Winders, T., & Greenhawt, M. (2020). Unmet needs of children with peanut allergy. *Annals of Allergy Asthma & Immunology*, 124(5), 479–486. <https://doi.org/10.1016/j.anai.2020.01.016>
- Cox, D. W., & Souëf, P. N. L. (2014). Human rhinovirus C and asthma in childhood. *Clinical Pulmonary Medicine*, 21(3), 107–112. <https://doi.org/10.1097/cpm.0000000000000036>
- Ralalage, D. D., & Hurst, J. R. (2023). Chronic obstructive pulmonary disease: aetiology, pathology, physiology and outcome. *Medicine*, 51(10), 737–741. <https://doi.org/10.1016/j.mpmed.2023.07.008>
- Needham, R. S., Sharpe, G. R., Williams, N. C., Lester, P. A., & Johnson, M. A. (2022). Exercise-induced bronchoconstriction in university field hockey athletes: Prevalence, sex differences, and associations with dyspnea symptoms. *Frontiers in Allergy*, 3, 994947. <https://doi.org/10.3389/falgy.2022.994947>
- Wu, X., Zhou, R., Zhong, Q., Wu, X.-Y., Chen, H., Wei, Y.-F., Xiong, L., & Wu, X.-X. (2025). Lung function, proteomic signatures, and risk of metabolic dysfunction-associated steatotic liver disease. <https://doi.org/10.21203/rs.3.rs-6381296/v1>

